



# The effects of nitrous oxide on behavior and physiological parameters during conscious sedation with a moderate dose of chloral hydrate and hydroxyzine

Wes McCann, DDS, MS   Stephen Wilson, DMD, MA, PhD   Peter Larsen, DDS   Bill Stehle, BS

## Abstract

The purpose of this study was to determine differences in heart rate (HR), blood pressure (BP), peripheral oxygen saturation ( $pO_2$ ), expired  $CO_2$  ( $CO_2$ ), and behavior (using two scales) comparing nitrous oxide/oxygen ( $N_2O$ ) with oxygen ( $O_2$ ) alone in 20 children (mean age  $45 \pm 5.1$  months) sedated with chloral hydrate (CH) and hydroxyzine in a double-blind crossover design. Administration of CH (40 mg/kg) and hydroxyzine (2 mg/kg) was held constant for each patient visit; however,  $N_2O$  (50%) was administered during one visit and  $O_2$  (100%) at the other in a randomly determined manner. Physiologic and behavioral parameters were collected during eight specific procedural events (e.g., administration of local anesthesia). Data were analyzed with a repeated-measures ANOVA, one-way ANOVA, t-test, Kruskal Wallis ANOVA, and descriptive statistics. There was no statistically significant difference in any physiologic or behavioral parameter as a function of inhalation agent. However, significant differences were found for certain physiological parameters (i.e., HR [ $F = 5.41$ ,  $P < 0.001$ ],  $pO_2$  [ $F = 6.04$ ,  $P < 0.001$ ], and  $CO_2$  [ $F = 2.33$ ,  $P < 0.027$ ]) and all behavioral measures (% crying [ $F = 2.82$ ,  $P < 0.008$ ], % quiet [ $F = 5.38$ ,  $P < 0.001$ ], % movement [ $F = 3.88$ ,  $P < 0.001$ ], and % struggle [ $F = 2.83$ ,  $P < 0.007$ ]) of one scale (Ohio State University Behavioral Rating Scale [OSUBRS]) as a function of procedural events. Although no statistically significant differences were attributable to inhalation agent, evidence suggests that  $N_2O$  resulted in less crying and struggling and more quiet behaviors than  $O_2$ . Significant correlations existed between sub-categories of the two behavioral rating scales, suggesting some association between the scales. One may conclude from the results of this study that moderate doses of CH and hydroxyzine in combination with nitrous oxide are not associated with any significant potentiation effects on physiologic parameters compared with the same oral agents with oxygen alone. Certain procedural events (e.g., administration of local anesthesia) do result

in patient responses that affect specific behaviors and physiology. Although the effects of  $N_2O$  may not be statistically significant, generally it produces an attenuation in physiological and behavioral responses as measured under the conditions of this study. (*Pediatr Dent* 18:35-41, 1996)

The most commonly used drug combination for the sedation of pediatric dental patients is CH and hydroxyzine supplemented with nitrous oxide ( $N_2O$ ).<sup>1,2</sup> A variety of studies have evaluated pediatric dental sedations utilizing various drugs or drug combinations supplemented with  $N_2O$ .<sup>3-13</sup> All used sedative agents in doses consistent with the recognized upper limits of therapeutic ranges, and none reported any significant adverse effects despite some probability of such.

Few studies have focused on the effects of  $N_2O$  on behavioral and physiological responses in young children sedated with a combination of common sedative agents used in a relatively moderate dose range. In theory, under most circumstances, the addition of  $N_2O$  to a combination of sedative agents should result in the reduced need for higher doses of the other agents, attaining the same sedation end-point because of the potentiation effect of  $N_2O$ . Concerns of  $N_2O$  as a potentiating agent during pediatric dental sedations have become widely discussed in light of the recently revised American Academy of Pediatrics guidelines concerning sedation procedures.<sup>14</sup>

Haupt et al.<sup>10</sup> studied the effects of  $N_2O$  on CH sedation in 20 children, (mean age of 32 months) by evaluating behavior at two appointments. Patients randomly received 50 mg/kg CH dose as a sedative agent with either 50%  $N_2O/O_2$  for the first 20 min of treatment then 100%  $O_2$  for the remainder of treatment; or 100%  $O_2$  for the first 20 minutes, switching to 50%  $N_2O/O_2$  for the remainder. The alternate inhalation agent regimen pairing was used during the second appointment. Results indicated improved sedative effects in about half of

patients when N<sub>2</sub>O was combined with CH. Behavior assessment involved a subjective rating scale for overall behavior during visits, which increased the likelihood of either a halo effect or a loss of procedurally related changes or both. Vital signs remained unchanged throughout treatment with the exception of slight pulse and respiratory rate elevations during episodes of increased oral stimulation (i.e., mouth prop insertion and local anesthetic administration). Similar physiologic responses have been noted in other studies.<sup>4, 5, 15-17</sup>

Shapira et al.<sup>11</sup> also reported behavior improvement in patients receiving N<sub>2</sub>O during sedation in a study comparing the effects of hydroxyzine alone, hydroxyzine with N<sub>2</sub>O, and N<sub>2</sub>O alone. However, the effects on physiology were not addressed. The limited knowledge of potentiating effects of N<sub>2</sub>O following administration of sedative agents warrants further investigation. To gain a better understanding of the effects of N<sub>2</sub>O on physiologic and behavioral responses during sedations of young children, this study had the following objectives:

1. To monitor and evaluate physiologic functions (HR, BP, CO<sub>2</sub> levels, and pO<sub>2</sub>) following a moderate dose of CH/hydroxyzine comparing N<sub>2</sub>O with O<sub>2</sub> alone
2. To evaluate the quality of sedations, based on operationally defined patient behavior, following CH/hydroxyzine administration comparing N<sub>2</sub>O with O<sub>2</sub> alone.

## Methodology

### Sample and design

Consent was obtained for 20 children (ages 36–60 months) to participate in this institutionally approved study. All were ASA class I, required more than one sedation visit for completion of operative dentistry, and exhibited a variety of the following behaviors during examination by a pediatric dentist or resident:

1. Failure to open mouth following directions to do so
2. Active attempts to escape from the dental chair
3. Extraneous flailing of arms or legs
4. Excessive interfering head movements.

A standardized dose of CH (40 mg/kg) and hydroxyzine (2 mg/kg) was used during each patient visit. N<sub>2</sub>O was administered at a ratio of 50% nitrous oxide and 50% oxygen.

Physiologic monitoring included: Critikon Dinamap Vital Signs Monitor (Dinamap™, Tampa, FL), 1846SX (BP); Nellcor Pulse Oximeter and Printer (Nellcor®, Hayward, CA), Model N-100 and N-9000, respectively (HR and pO<sub>2</sub>); Datex Carbon Dioxide Monitor (Datex™, Helsinki, Finland), Model 223 (CO<sub>2</sub>). The Porter MXR nitrous oxide delivery system was used. Monitoring of behavioral patterns was carried out by vid-

eotaping each treatment session and analyzing the tapes later.

Patients were assigned randomly to one of two groups (A or B) in a double-blind cross-over design. At the first appointment, group A received CH (40 mg/kg) and hydroxyzine (2 mg/kg) PO, was observed for a 45-min latency period, then treated with N<sub>2</sub>O via the nasal hood. Group B participants received equivalent dosages of CH and hydroxyzine PO, were also observed for a 45-min latency period, and were treated with 100% O<sub>2</sub> administration via the nasal hood. The alternate regimen was administered at the second appointment. Scheduled appointments were at the same time of day (8–12 am) and were no more than 4 weeks apart.

### Preoperative steps

Baseline physiologic values (BP, HR, pO<sub>2</sub>, and CO<sub>2</sub> levels) were obtained. The BP cuff was placed on the right arm, and the HR and pO<sub>2</sub> probe was affixed to the right index finger during baseline and the right middle toe during treatment. Baseline CO<sub>2</sub> values were obtained from a cannula in the right naris and recordings from patient's normal respiratory pattern. Treatment values were obtained similarly irrespective of regimen.

Parent or operator administered medications mixed with syrup and carbonated beverage. Parents and patient waited in the reception room for 45 min then a secondary operator escorted the child to the treatment room without the primary operator and started videotaping. Either N<sub>2</sub>O (titrated initially, then to 50% at 5 L/min) or O<sub>2</sub> (100% at 5 L/min), depending on the child's assigned group, was started using tell-show-do and positive reinforcement. During this initial titration phase, the need for a Papoose Board™ (Olympic Medical Group, Seattle, WA) was determined based on the patient exhibiting one of the following three behaviors: 1) excessive struggling/fighting without use of voice control; 2) high hands and attempts to escape from the chair; or 3) arm and/or leg flailing/kicking that did not cease following tell-show-do or voice control requests. Patients were not wrapped in the Papoose Board unless behavior warranted it. The N<sub>2</sub>O delivery system was placed out of view of the primary operator who returned to the treatment room and initiated treatment once the patient was stabilized.

Physiological parameters were recorded throughout the treatment session by automated monitor recorders and manually by another assistant. Operative or extraction procedures were limited to posterior quadrants and lasted approximately 60 min.

### Data analysis

Behavior was rated and analyzed later based on the following eight segments of each session:

1. Start of videotape until topical application
2. Topical anesthetic application
3. Local anesthetic injection

4. Rubber dam placement
5. Initiation of operative procedures
6. 5 min following treatment initiation
7. 10 min following treatment initiation
8. 15 min following treatment initiation.

The study used the Ohio State University Behavior Rating Scale (OSUBRS), which involves four behavioral categories based on head or bodily movements, crying and oral/physical resistance as follows:

1. Q = quiet behavior, no movement
2. C = crying with no struggling
3. M = movement with struggling only, no crying
4. S = crying and struggling exhibited simultaneously, disruptive behavior. Struggling is defined as rapid or intense head, foot, or hand movements.

The Automated Counting System (ACS)<sup>TM</sup> (Version 1.0 JAGTECH, Rockville, MD) computer software program, was used to quantify behavioral categories by the rater who recorded each behavioral category by pressing its respective keyboard key (i.e., Q, C, M, S). Any change from one behavioral category to another was noted by pressing the appropriate key. Behavioral categories were mutually exclusive and only one was identified for any given time period. A computerized printout of data provided information on the frequency, duration, and mean duration of each category of behavior during any defined segment of the treatment session.

Previous studies indicated that intra- and inter-rater reliability as measured by a correlation analysis was 95–99%.<sup>19,20</sup> Intrarater reliability was measured by a random selection of three treatment visits, which were initially rated and, following the 20th rating session, rated again. Additionally, behaviors were rated intraoperatively by the operator or assistant using a simple ordinal scale that categorized behavior as either quiet, struggling, or crying. For statistical purposes, the following scores were assigned to the behavioral categories: 1 = quiet; 2 = struggling; and 3 = crying. These ratings occurred at every procedural event identified previously and referred only to the observed behavior at the time of the event.

The mean, standard deviation, and frequency distribution were used to characterize the age and sex of the sample population. Intrarater reliability was evaluated by *t*-test analysis comparing the first to the second rating for each behavioral category for the OSUBRS. A repeated-measures ANOVA was used to determine individual significant changes in the OSUBRS categories and physiological parameters across the eight procedural events as a function of inhalation agent (N<sub>2</sub>O versus O<sub>2</sub>). The nonparametric Kruskal-Wallis one-way ANOVA was utilized to analyze behavior ratings associated with the intraoperative ordinal scale. Finally, a correlation coefficient was used to determine any sig-

nificant association between the two behavioral rating scales used across the procedural events.

## Results

Physiologic and behavioral data were collected from 40 sedation visits involving 26 males and 14 females, ages 36 to 55 months (mean = 45 months). Weight ranged from 13.0 to 20.5 kg (mean = 16.07 kg). Utilizing a set dosage per kg body weight for chloral hydrate (40 mg/kg) and hydroxyzine (2 mg/kg), the mean dosage administered was 637 mg and 32 mg, respectively.

### Physiologic response

A repeated-measures ANOVA revealed no statistically significant differences in any physiologic parameter as a function of inhalation agent; however, significant differences for procedural events were found for HR ( $F = 5.41, P < 0.001$ ), pO<sub>2</sub> ( $F = 6.04, P < 0.001$ ), and CO<sub>2</sub> ( $F = 2.33, P < 0.027$ , see [Table 1]). Although no significant difference was found comparing inhalation agents, Figs 1 and 2 demonstrate generally that N<sub>2</sub>O tended to have an attenuating effect compared with O<sub>2</sub> on mean HR and CO<sub>2</sub> (actually CO<sub>2</sub> was within normal limits compared with pO<sub>2</sub>, which decreased due to crying), although not as consistent as the behavioral data.


### Behavioral response

Behavior categories of quiet (Q), crying (C), struggling (S), and movement (M) were quantified by a single rater viewing videotapes of each sedation appointment. A *t*-test comparing first and second reliability sessions for each behavioral category revealed no significant difference ( $P = 0.05$ ).


**TABLE 1. REPEATED MEASURES ANOVA OF PHYSIOLOGIC PARAMETERS AS A FUNCTION OF PROCEDURE AND INHALATION AGENT**

Parameter	Variable	df	F-Value	P
Heart rate	N <sub>2</sub> O	1	0.64	0.43
	Procedure	7	5.41	0.001*
	N <sub>2</sub> O x Proc	7	0.63	0.63
Systolic BP	N <sub>2</sub> O	1	0.09	0.76
	Procedure	7	1.77	0.093
	N <sub>2</sub> O x Proc	7	0.078	0.078
Diastolic BP	N <sub>2</sub> O	1	0.15	0.70
	Procedure	7	1.11	0.356
	N <sub>2</sub> O x Proc	7	1.12	0.353
O <sub>2</sub> sat	N <sub>2</sub> O	1	0.15	0.70
	Procedure	7	6.04	0.001*
	N <sub>2</sub> O x Proc	7	0.72	0.658
Expired CO <sub>2</sub>	N <sub>2</sub> O	1	0.58	0.45
	Procedure	7	2.33	0.027*
	N <sub>2</sub> O x Proc	7	0.98	0.44

\* Significant.



**Fig 1. Mean heart rate across procedures as a function of inhalation agent.**



**Fig 2. Mean expired CO<sub>2</sub> across procedures as a function of inhalation agent.**

**TABLE 2. MEAN (± SD) PERCENT BEHAVIORS OF OSUBRS\* ACROSS PROCEDURES BY INHALATION AGENTS**

Behavior	Procedure															
	Baseline		Topical		Local		Rubber Dam		Start		5 Min		10 Min		15 Min	
	N <sub>2</sub> O	O <sub>2</sub>	N <sub>2</sub> O	O <sub>2</sub>	N <sub>2</sub> O	O <sub>2</sub>	N <sub>2</sub> O	O <sub>2</sub>	N <sub>2</sub> O	O <sub>2</sub>	N <sub>2</sub> O	O <sub>2</sub>	N <sub>2</sub> O	O <sub>2</sub>	N <sub>2</sub> O	O <sub>2</sub>
% Cry	8.3 <sup>†</sup>	18.8	12.6	14.6	28.1	33.3	12.6	20.1	19.5	14.5	13.3	24.0	11.7	10.4	4.7	21.6
	18.0 <sup>†</sup>	25.6	28.9	31.6	31.3	31.8	29.8	35.9	35.0	30.0	28.1	35.9	27.4	25.9	13.4	34.4
% Movement	4.8	5.7	2.0	2.8	4.1	2.7	3.7	1.5	0.2	0.3	2.4	0.5	1.4	0.3	0.4	1.69
	6.7	6.0	6.5	9.9	8.8	5.0	9.8	5.0	0.9	1.6	5.8	2.1	4.5	1.4	1.1	4.3
% Quiet	85.1	70.3	81.9	72.6	51.7	47.6	74.0	72.2	75.2	78.5	82.4	66.8	83.2	84.6	94.5	73.7
	20.3	29.8	36.5	39.6	40.1	39.6	41.8	43.6	41.5	39.9	31.5	46.4	34.3	36.5	14.2	38.3
% Struggle	1.6	5.0	3.4	9.8	15.9	16.2	9.61	6.0	4.9	6.5	1.8	8.5	3.5	4.5	0.23	2.9
	5.1	8.5	10.5	29.5	26.0	24.0	25.4	17.2	16.7	15.9	5.1	18.0	9.75	11.9	1.0	6.4

\* Ohio State University Behavioral Rating Scale.

<sup>†</sup> Mean.

<sup>‡</sup> SD.

Percent duration of each behavioral category of the OSUBRS was calculated. A repeated-measures ANOVA indicated that there were no statistically significant differences in any behavioral category as a function of N<sub>2</sub>O versus O<sub>2</sub>. However, significant differences for procedural events were found for all categories (%C [F = 2.82, P < 0.008], %Q [F = 5.38, P < 0.001], %M [F = 3.88, P < 0.001], and %S [F = 2.83, P < 0.007]). Table 2 shows a summary of the mean occurrence of each behavioral category over the rated procedural periods. Of note is the wide variability observed across all categories of behavior, especially for infrequently occurring behaviors (M and S).

Intraoperative behaviors were recorded, coincidental to physiological data, based on the following scale: 1 = Q; 2 = S; 3 = C (the scale was not evaluated for reliability). From these data, Kruskal Wallis one-way ANOVA revealed no statistically significant differences in behavior with respect to inhalation agent. However, Fig 3 demonstrates a lower mean score for the N<sub>2</sub>O visit interpreted as more quiet behavior compared with O<sub>2</sub>. There was a statistically significant and modest association between the categories of the OSUBRS and the simple clinical scale. Table 3 shows the correlation co-

efficients. The Papoose Board was required for 10 visits when N<sub>2</sub>O was used and for 14 visits with O<sub>2</sub>.

## Discussion

We found no statistical difference between N<sub>2</sub>O and O<sub>2</sub> conditions for the physiologic and behavioral parameters measured. Graphically, N<sub>2</sub>O appeared to limit behavioral responsiveness and some physiological parameters and this trend may contribute to a common conviction that N<sub>2</sub>O has beneficial effects during sedation. Certain physiological parameters and all behavioral categories of the OSUBRS were influenced significantly by procedural events.

## Physiology

Although no statistical difference in physiological parameters as a function of inhalation agent was detected, a consistent trend was observed graphically for mean HR and CO<sub>2</sub>. Data from this study suggest that N<sub>2</sub>O decreases mean HR, and that pO<sub>2</sub> decreases mean CO<sub>2</sub> (due to crying) in response to procedural events. It is not likely that these trends were due directly to the pharmacologic action of N<sub>2</sub>O alone. A slight depressing effect on myocardial contractility has been reported with N<sub>2</sub>O, but N<sub>2</sub>O in a therapeutic range used in den-


**TABLE 3. CORRELATION COEFFICIENTS SHOW MODEST ASSOCIATION BETWEEN BEHAVIORAL CATEGORIES OF THE OSUBRS\* AND THE SIMPLE CLINICAL SCALE**

Behaviors	Baseline	Topical	Local	Rubber Dam	Start	5 Min	10 min	15 min
Crying	0.58 <sup>†</sup>	0.67 <sup>†</sup>	0.71 <sup>†</sup>	0.82 <sup>†</sup>	0.82 <sup>†</sup>	0.34	0.73 <sup>†</sup>	0.53 <sup>‡</sup>
Struggle	0.21	0.39	0.36	0.71 <sup>†</sup>	0.55 <sup>†</sup>	0.33	0.66 <sup>†</sup>	0.40
Quiet	-0.57 <sup>†</sup>	-0.74 <sup>†</sup>	-0.74 <sup>†</sup>	-0.96 <sup>†</sup>	-0.84 <sup>†</sup>	-0.33	-0.73 <sup>†</sup>	-0.52 <sup>†</sup>

\* Ohio State University Behavioral Rating Scale.

<sup>†</sup>  $P < 0.001$ .

<sup>‡</sup>  $P < 0.01$ .



**Fig 3. Mean behavior scores of simple clinical scale across procedures as a function of inhalation agent.**

istry has no known effect on HR or cardiac output.<sup>21</sup> More than likely, the observed mean physiologic parameters during N<sub>2</sub>O conditions would not be attributed to direct cardiovascular depression, but to greater patient relaxation.

Minimal variability existed for pO<sub>2</sub> values and no desaturation episodes of clinical consequence (< 95%)<sup>13, 22</sup> were observed without simultaneous foot/toe movements in this study. Procedural effects did cause significant but clinically irrelevant fluctuations in pO<sub>2</sub>. Patient movement has been shown to be a cause of false peripheral oxygen desaturations as measured by pulse oximetry.<sup>23, 24</sup> Because pO<sub>2</sub> normally demonstrates very little variability, minor fluctuations associated with conditions unrelated to pharmacological influence (e.g., Valsalva's maneuver and crying) can lead to erroneous and inappropriate conclusions.

Various studies have implicated advantages of sidestream capnography during pediatric dental sedations.<sup>25-28</sup> These advantages focus on the ability of capnography to detect apnea, airway obstruction and developing hypoventilation.<sup>28, 29</sup> Classical waveform patterns with normal respiratory rates are more typically observed in the quiet or sleeping patient. Changes in the waveform pattern can reflect many distinct conditions and require vigilance in monitoring. Any patient displaying waveform deviations was immediately postured with the head-tilt/chin-lift maneuver and evaluated with readjustments of the sampling port location in the naris or re-assessment of patient status. These precautions, along with 50-100% oxygen supple-

mentation, may have precluded desaturation episodes detected by pulse oximetry.

Slight differences in trends of CO<sub>2</sub> levels may be attributed to differences in clinical behavior (i.e., decreased means associated with increased

crying bouts) rather than to pharmacologic-induced conditions. In this study, patterns of CO<sub>2</sub> levels in both N<sub>2</sub>O and O<sub>2</sub> groups coincided with other physiological parameters measured and patient behavior, suggesting the latter's influence rather than a pharmacologic effect. These findings and interpretation support the findings of others.<sup>17, 25</sup>

HR and CO<sub>2</sub> monitoring are expected to best reflect children's responsiveness to procedures based on characteristics associated with stressful experiences. Salient stimuli will result in increased HR, the common mode of cardiovascular response to perceived stressful conditions in young children. Similar trends in clinical behavior have been reported by Houpt et al.<sup>10</sup> Likewise, under such conditions, children also respond by crying, resulting in decreased CO<sub>2</sub> concentration due to oral shunting of expired air.

### Behavior

Compared with O<sub>2</sub> alone, improved clinical behavior was observed during N<sub>2</sub>O visits in the categories of crying, quiet, and struggling. Movement occurred infrequently and was essentially equivalent for both agents. Fewer patients required restraint in the Papoose Board with N<sub>2</sub>O compared with O<sub>2</sub>. Increased Q and decreased C behaviors in the majority of patients occurred when N<sub>2</sub>O was used. From a clinical standpoint, the results indicate N<sub>2</sub>O used in combination with a modest dose of CH and hydroxyzine may improve intraoperative behavior for patients 36-60 months of age without significantly affecting physiologic function.

In a dose-response study, Wilson<sup>17</sup> found that during low doses of CH, accentuated physiologic responses notably occurred during local anesthetic injection and/or tooth preparation, but responses were dampened as doses increased. This observation was attributed to a deeper level of sedation imparted by increased CH dosage. He noted that significantly high doses of CH would be necessary to overcome more discomforting stimuli of certain dental procedures in many young children and that such a practice may lead to deep sedation, compromising patient safety.

Results of this study indirectly suggest that improved responses to procedures may not require significantly higher doses of CH. Although no statistical difference was detected as a result of inhalation agent, definitive clinical conclusions should be cautioned because of wide variation in patient responses (i.e., large standard deviations of the mean percent of each behav-

ioral category of the OSUBRS) and operator expectations. This study concurs with findings by Houpt et al.,<sup>10</sup> which imply that N<sub>2</sub>O does improve behavioral responses to stimuli in some individuals (viz., sedation augmentation was not consistently observed among subjects). Larger sample sizes, more refined patient selection techniques, and more sensitive behavioral and physiologic measures might demonstrate both clinical and statistically significant differences.

Although no scales were used in this study to rank sedation in terms of success/failure, the high percentage of quiet behavior (compared with crying, struggling, and movement) is an indirect indication that sedation efforts were effective, regardless of inhalation agent used. Treatment planned for each visit was completed without compromise; no physiologic aberrations were observed; and no vomiting occurred intraoperatively. Furthermore, no postoperative complications were noted or reported by parent/guardian.

One interesting finding was the association between the OSUBRS and a clinically relevant and simple technique of rating behavior. Correlations between the two scales involving Q and C categories were high and modest with S as a function of procedural events. Since the OSUBRS requires the tedious and costly procedure of videotaping and playback review, a simple technique such as rating behavioral response for each procedure occurring as either quiet, crying, or struggling may be economically advantageous and more feasible. It is not clear how much information is "lost" because only a brief moment in time is rated for the simple clinical rating scale, whereas the OSUBRS collects raw data continuously. It's analogous to the situation of comparing the output of an automated BP cuff set to record BP and HR every 5 min with that of continual output of HR and pO<sub>2</sub> of a pulse oximeter. In the former, a significant amount of information is not retrievable. Further study into this issue is warranted.

## Conclusions

In pediatric dental sedations, utilizing 40 mg/kg CH plus 2 mg/kg hydroxyzine:

1. No physiologic parameters were significantly affected by the addition of inhalation agents to a dose of chloral hydrate and hydroxyzine in the moderate therapeutic range
2. HR, pO<sub>2</sub>, and CO<sub>2</sub> were affected significantly by dental procedures. HR increased and CO<sub>2</sub> decreased reflecting patient discomfort and crying, particularly during procedural phases of injection and tooth preparation
3. OSUBRS categories of percent crying increased during O<sub>2</sub> visits while percent quiet increased during N<sub>2</sub>O visits. A simple clinical rating scale correlated well with the OSUBRS and reflected similar changes.

We extend sincere appreciation to Deborah Hines for her patience, understanding, and efficient assistance throughout the course of this study.

Dr. McCann is in private practice in Memphis Tennessee. Dr. Wilson is Associate Professor and Director, Pediatric Dentistry Residency Program, The Ohio State University and Columbus Children's Hospital, Columbus, Ohio. Dr. Larsen is Associate Professor and Director, Oral and Maxillofacial Surgery Residency Program, and Dr. Stehle is a resident in periodontology, both at The Ohio State University, College of Dentistry, Columbus.

1. Houpt MI: Report of project USAP: the use of sedative agents in pediatric dentistry. *ASDC J Dent Child* 56:302-9, 1989.
2. Houpt MI: Project USAP the use of sedative agents in pediatric dentistry: 1991 update. *Pediatr Dent* 15:36-40, 1993.
3. Houpt MI, Sheskin RB, Koenigsberg SR, Dejardins P, Shey Z: Assessing chloral hydrate dosage for young children. *ASDC J Dent Child* 52:364-69, 1985.
4. Koenigsberg SR, Houpt MI, Weiss NJ: A comparison of chloral hydrate with and without promethazine in the premedication of children. *J Dent Res* 63:233, 1984.
5. Houpt MI, Koenigsberg SR, Weiss NJ, Desjardins PJ: A comparison of chloral hydrate with and without promethazine in the sedation of young children. *Pediatr Dent* 7:41-46, 1985.
6. Poorman TL, Farrington FH, Mourino AP: Comparison of a chloral hydrate/hydroxyzine combination with and without meperidine in the sedation of pediatric dental patients. *Pediatr Dent* 12:288-91, 1990.
7. Meyer ML, Mourino AP, Farrington FH: Comparison of triazolam to chloral hydrate/hydroxyzine combination in the sedation of pediatric dental patients. *Pediatr Dent* 12:283-87, 1990.
8. Badalaty MM, Houpt MI, Koenigsberg SR, Maxwell KC, DesJardins PJ: A comparison of chloral hydrate and diazepam sedation in young children. *Pediatr Dent* 12:33-37, 1990.
9. Moody EH, Mourino AP, Campbell RL: The therapeutic effectiveness of nitrous oxide and chloral hydrate administered orally, rectally, and combined with hydroxyzine for pediatric dentistry. *ASDC J Dent Child* 53:425-29, 1986.
10. Houpt MI, Rosivack RG, Rozenfarb N, Koenigsberg SR: Effects of nitrous oxide on chloral hydrate sedation of young children. *Anesth Prog* 33:298-302, 1986.
11. Shapira J, Holan G, Guelmann M, Cahan S: Evaluation of the effect of nitrous oxide and hydroxyzine in controlling the behavior of the pediatric dental patient. *Pediatr Dent* 14:167-70, 1992.
12. Houpt MI, Manetas C, Joshi A, DesJardins PJ: Effects of chloral hydrate on nitrous oxide sedation of children. *Pediatr Dent* 11:26-29, 1989.
13. Mueller WA, Drummond JN, Pribisco TA, and Kaplan RF: Pulse oximetry monitoring of sedated pediatric dental patients. *Anesth Prog* 32:237-40, 1985.
14. American Academy of Pediatrics, Committee on Drugs: Guidelines for Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures. *Pediatrics* 89:1110-15, 1992.
15. Doring KR: Evaluation of an alphaprodine-hydroxyzine combination as a sedative agent in the treatment of the pediatric patient. *J Am Dent Assoc* 111:567-76, 1985.
16. Currie WR, Biery KA, Campbell RL, Mourino AP: Narcotic sedation and evaluation of cardiopulmonary parameters and behavior modification in pediatric dental patients. *J Pedod* 12:230-49, 1988.
17. Wilson S: Chloral hydrate and its effects on multiple physiological parameters in young children: a dose-response study. *Pediatr Dent* 14:171-77, 1992.

18. Chambers WL, Fields HW, Machen JB: Measuring selected disruptive behaviors of the 30- to 60-month-old patient. Part 1: Development and assessment of a rating scale. *Pediatr Dent* 3:251-56, 1981.
19. Lochary ME, Wilson S, Griffen AL, Coury DL: Temperament as a predictor of behavior for conscious sedation in dentistry. *Pediatr Dent* 15:348-52, 1993.
20. Radis FG, Wilson S, Griffen AL, Coury DL: Temperament as a predictor of behavior during initial dental examination in children. *Pediatr Dent* 16:121-27, 1994.
21. Malamed SF: Sedation: a guide to patient management. St Louis: CV Mosby Co, 1989, p 102.
22. Whitehead BG, Durr DP, Adair SM, Proskin HM: Monitoring of sedated pediatric dental patients. *ASDC J Dent Child* 55:329-33, 1988.
23. Wilson S: Conscious sedation and pulse oximetry: false alarms? *Pediatr Dent* 12:228-32, 1990.
24. Verwest TM, Primosch RE, Courts FJ: Variables influencing hemoglobin oxygen desaturation in children during routine restorative dentistry. *Pediatr Dent* 15:25-29, 1993.
25. Iwasaki J, Vann WF Jr, Dilley DCH, Anderson JA: An investigation of capnography and pulse oximetry as monitors of pediatric patients sedated for dental treatment. *Pediatr Dent* 11:111-17, 1989.
26. Wilson S, Tafaro ST, Vieth RF: Electromyography: its potential as an adjunct to other monitored parameters during conscious sedation in children receiving dental treatment. *Anesth Prog* 37:11-15, 1990.
27. Anderson JA: Respiratory monitoring for anesthesia and sedation. *Anesth Prog* 34:228-31, 1987.
28. Anderson JA, Vann WF Jr: Respiratory monitoring during pediatric sedation: pulse oximetry and capnography. *Pediatr Dent* 10:94-101, 1988.
29. Wilson S: Patient monitoring in the conscious sedation of children for dental care. *Curr Opin Dent* 1:570-76, 1991.

## Earn CE credits while you read this journal



Earn credits from your home on your own time. No need to travel; no long lectures to attend. *Pediatric Dentistry* now offers up to 18 continuing education credits per year for demonstrating an understanding of topics discussed in selected journal articles.

It couldn't be easier. As a subscriber, you will receive a multiple-choice test covering several articles around the same time you receive the journal. Simply read the selected articles and return your answer sheets to AAPD for grading. We will notify you of the number of credits you earned for your correct answers.

The CE logos on the cover of the journal and on the title pages indicate which articles will be tested. Annual subscription price is \$60.

A related service, the Continuing Education Registry, helps you keep track of your CE credits. Subscribers will receive reporting forms on which to record continuing education credits. For \$30 per year, credit information submitted to AAPD will be entered into a confidential record. Reports will be furnished annually or by request. If you subscribe to both services, your journal CE credits will be entered automatically into the CE Registry. See page 2 to subscribe to both new member services.