

## Effects of formocresol alone vs. formocresol with eugenol on macrophage adhesion to plastic surfaces

Juan José Segura, DDS, MD, PhD Alicia Jiménez-Rubio, DDS, MD, PhD Juan Ramón Calvo, MD, PhD

### Abstract

**Purpose:** *The purpose of this study was to compare the in-vitro effects of a European-based formocresol formulation that incorporates eugenol with formocresol alone on the adhesion of macrophages to plastic surfaces.*

**Methods:** *Macrophages were obtained from Wistar™ rats. The adherence capacity of macrophages to a plastic surface was determined. Assays were carried out in Eppendorf® tubes incubated for 15 min at 37°C in a humidified atmosphere of 5% CO<sub>2</sub>. The adherence index was calculated.*

**Results:** *Results showed that both formocresol/eugenol and formocresol alone significantly decreased the adherence index of macrophages. The formocresol formulation that incorporated eugenol was more potent in inhibiting macrophage adhesion than formocresol alone.*

**Conclusions:** *Taking into account that adherence to a substrate is the first step in the phagocytic process of macrophages and in antigen presentation, both formocresol formulations could inhibit macrophage function and modulate immune and inflammatory responses in dental pulp and periapical tissues (Pediatr Dent 20:3 177–80, 1998).*

Formocresol (Buckley's formula) consists of 19% formaldehyde and 35% cresol in a vehicle of 15% glycerin in water.<sup>1</sup> This formocresol formulation, as well as modifications that contain different proportions of formaldehyde and cresol, are widely used by American dentists for pulpotomy procedures in primary teeth<sup>2,3</sup> and as an endodontic drug to neutralize the septic and necrotic contents of root canal.<sup>4,5</sup>

Formocresol is highly toxic to cells.<sup>6,7</sup> Tissue changes in dog livers and kidneys induced by the absorption of formocresol from pulpotomy sites have been shown.<sup>8</sup> Polyvinyl sponge implants containing full-strength formocresol cause fixation of fibroblasts in adjacent cells.<sup>9</sup> A 1:5 dilution of formocresol markedly suppressed lactic dehydrogenase activity.<sup>10</sup> At weaker concentrations, formocresol does not fix the tissue, but might create signs of cellular degeneration.<sup>11</sup> Human clinical studies have shown that formocresol treatment causes severe inflammatory reactions or necrosis of the pulp.<sup>12</sup>

However, the amount of formocresol that could be absorbed from pulpotomy sites is minimal.<sup>13</sup> In the light of the preceding discussion, the use of formaldehyde-containing pastes on pulp tissues is controversial.<sup>5,9</sup>

In Europe, therapeutic pulpotomies of primary teeth are performed using a formulation of formocresol which incorporates eugenol (20% formaldehyde, 20% tricresol, and 20% eugenol). Eugenol is an ingredient in many over-the-counter toothache drops, temporary filling materials, restorative materials, bases, cements, intracanal medicaments, and in pastes used in endodontics as a sealer for root canal fillings.<sup>14–16</sup> It has been claimed that eugenol does not irritate the pulp if applied to dentin. In contrast, Glass and Zander<sup>17</sup> showed that eugenol induced chronic inflammation when placed on an exposed pulp. Moreover, Brännström and Nyborg<sup>18</sup> and Webb and Bussell<sup>19</sup> also demonstrated that eugenol produces a marked inflammatory reaction in pulp.

On the other hand, it has been shown that inflamed pulp and periapical tissues contain a variety of immunocompetent cells, with macrophages predominating.<sup>20</sup> Moreover, macrophages are the most dominant immunocompetent cells through all stages of induced periapical lesions.<sup>21,22</sup> Macrophages are implicated in chronic inflammation and repair of pulpal and periapical tissues.<sup>23</sup> They are known to have several mediator and regulatory functions, and are involved in the entire spectrum of the defense reactions.<sup>24</sup> It is well documented that adherence is the first step in the phagocytic process of inflammatory macrophages.<sup>25</sup>

The in-vitro effects of formocresol alone and a formocresol formulation that incorporates eugenol on the substrate adherence capacity of rat inflammatory macrophages are compared in this study.

### Methods

Formocresol/eugenol (20% formaldehyde, 20% tricresol, and 20% eugenol) was obtained from Laboratono Bucca (Juan Alvarez Mendizabal, Madrid). Formocresol (42.1% cresol, 42.1% formaldehyde, 15.8% ethilic alcohol) was obtained from J. Bird

Moyer. RPMI-1640 medium was obtained from Sigma (St. Louis, MO).

The protocol was approved by our experimentation committee. Peritoneal macrophages were elicited from Wistar rats.<sup>26</sup> Briefly, each rat was injected intraperitoneally with 5 mL of sterile 6% sodium caseinate. Animals were killed after 4 days by decapitation and the peritoneal cavity was washed with 10 mL of cold 0.9% NaCl. After a 2-min massage, the cell exudate was removed with a syringe and centrifuged for 10 min at 250 x g at 4°C. The contaminating red blood cells were lysed with cold 0.2% NaCl. The remaining cells were then washed with 0.9% NaCl by centrifugation, resuspended in RPMI-1640 medium, counted, adjusted in the same medium at 2–4 x 10<sup>6</sup> macrophages/mL and immediately used for experiments. Mean cells per rat varied from 20–30 x 10<sup>6</sup>, of which 85–95% were macrophages by morphological criteria in Giemsa and Papanicolaou staining techniques. Viability, as determined by trypan-blue exclusion, was always greater than 94%.

The quantification of substrate-adherence capacity was carried out according to the technique described previously by De la Fuente et al.<sup>27</sup> with minor modifications. Aliquots of 180 µL of cell suspension were dispensed in Eppendorf tubes, which mimic the adher-

ence characteristics to tissues as reported by Noga et al.<sup>28</sup> Formocresol alone or formocresol/eugenol (20 µL) was dissolved directly in RPMI-1640 medium to a final dilution of 1:10, 1:100, or 1:1000 in the incubation medium. RPMI-1640 medium (20 µL) was added to control samples. Adherence assays were performed with 15 min of incubation at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> to provide a maximal adherence index.<sup>21</sup> After gentle centrifugation to remove nonadherent cells (3 s in the vortex in position 5), 10-µL aliquots were taken from each sample and the number of nonadherent macrophages/mL were counted in Neubauer chambers. No agglutination of macrophages was observed. The adherence index (AI) was calculated according to the following equation:

$$AI = 100 - \frac{\text{Nonadherent macrophages/mL}}{\text{Initial macrophages/mL}} \times 100$$

All values were expressed as the mean ± standard deviation of five separate experiments performed in triplicate, as indicated in the corresponding figure. The data were statistically evaluated by ANOVA. A value of *P* < 0.05 was considered statistically significant.

## Results

Both formocresol alone and formocresol/eugenol inhibited the substrate-adherence capacity of macrophages in a dose-dependent manner (Fig 1). When added to the incubation medium at a final dilution of 1:1000, formocresol/eugenol decreased the AI by 12% (*P* < 0.05). Moreover, the 1:100 dilution of formocresol/eugenol significantly decreased the AI by 59% (*P* < 0.05). The 1:10 dilution of formocresol/eugenol strongly and very significantly decreased the AI by 94% (*P* < 0.01). Inhibition (half of the maximal) of AI (IC<sub>50</sub>) was obtained at 1:327.3 formocresol/eugenol dilution.

Formocresol alone at a final dilution of 1:1000 did not significantly decrease the AI of macrophages (*P* > 0.05). However, the 1:100 dilution of formocresol significantly decreased the AI by 26% (*P* < 0.05). The 1:10 dilution of formocresol decreased the AI of macrophages by 78% (*P* < 0.05). Inhibition (half of the maximal) of AI (IC<sub>50</sub>) was obtained at 1:48.5 formocresol dilution.

## Discussion

In our study we demonstrate that both formocresol alone and formocresol/eugenol decrease the in-vitro substrate-adherence capacity of rat peritoneal macrophages to plastics surfaces. Although it has been shown that the adherence of macrophages to plastic requires cell surface characteristics different from adherence to collagen, fibronectin, or other cells,<sup>29</sup> the macrophage adhesion to a smooth plastic surface is comparable to that taking place in animal tissues.<sup>26–28, 30, 31</sup>

The sensitivity of cells to formocresol alone and formocresol/eugenol dilutions as high as 1:100, which




Fig 1. Dilution-effect curve for the effects of formocresol alone and formocresol/eugenol on AI of rat peritoneal macrophages. Macrophages (2–4 x 10<sup>6</sup>/inl) were incubated at 37°C in humidified, 5% CO<sub>2</sub> in the absence (C) or presence of decreasing dilutions (increasing concentrations) of formocresol alone (C) and formocresol/eugenol (e). After 15 min reaction was stopped and the AI calculated. Each point is the mean of five experiments performed in triplicate. \* *P* < 0.05 vs. control (C); \*\* *P* < 0.01 vs. control (C).

are similar to those found in periapical tissues, suggests that their inhibitory effect on macrophage adhesion may have physiological significance at the level of periapical tissues following formocresol/eugenol pulpotomy if this substance leaks through the apical foramen and invades the periapical zone.

Formocresol alone was less potent in inhibiting macrophage adhesion than formocresol formulations containing eugenol. The increased inhibitory effect of formocresol/eugenol on the substrate-adherence capacity of macrophages could be due to the chelating effect of eugenol on calcium ions.<sup>32</sup> The substrate adherence capacity of macrophages requires the presence of calcium ions in the medium.<sup>33</sup> Thus, the chelating action of eugenol on calcium ions could decrease macrophage adhesion.

## Conclusions

Formocresol alone is widely used by American clinicians for pulpotomy procedures.<sup>2</sup> In Europe, on the contrary, a formocresol formulation incorporating eugenol is frequently used. Our results demonstrate that formocresol alone is less toxic to macrophage adhesion than formocresol/eugenol.

Macrophages play an essential role in the immune response of the host to inflammatory and infectious processes, as well as in the reparative process. At the periapical-tissue level, macrophages, with phagocytosis and antigen presentation, have a central function in the repair of chronic apical periodontitis.<sup>21, 23</sup>

The influx of macrophages into the inflamed periapical tissues was most evident between 0 and 3 days after the pulp exposure.<sup>21</sup> The periapical tissues are highly responsive to pulpal injury, and begin to work rapidly as a second line of local defense to eliminate noxious stimuli invading pulp.<sup>22</sup> Then, when formocresol, alone or combined with eugenol, is used in pulpotomy procedures or during endodontic therapy, macrophage function could be modified and the reparative mechanisms and inflammatory reactions could be altered at the pulpal and periapical-tissue level.

This investigation was supported by grants from Dirección General de Investigación Científica y Técnica (DGICYT, PB94-1434 and PM9S-0159), Ministry of Education and Culture of Spain.

Dr. Segura is associate professor in the Departments of Dental Pathology and Therapeutics in the School of Dentistry and Medical Biochemistry and Molecular Biology in the School of Medicine, Dr. Jiménez-Rubio is associate professor in the Department of Dental Pathology and Therapeutics, School of Dentistry, and Dr. Calvo is lecturer in the Department of Medical Biochemistry and Molecular Biology, School of Medicine and Dentistry, all at the University of Seville, Seville, Spain.

## References

1. American Dental Association: Accepted dental therapeutics, 40th ed. Chicago, 1984.

2. McDonald RE, Avery DR: Tratamiento de las caries profundas, de la exposición pulpar y de los dientes desulpados. In *Odontología Pediátrica y del Adolescente*, 5th Ed. McDonald RE, Avery DR, Eds. Buenos Aires: Editorial Médica Panamericana SA, pp 408-435, 1990.
3. Morawa AP, Stratton LH, Han SS, Copron RE: Clinical evaluation of pulpotomies using dilute formocresol. *ASDC J Dent Child* 42:360-63, 1975.
4. Wesley DJ, Marshall FJ, Rosen S: The quantitation of formocresol as a root canal medicament. *Oral Surg Oral Med Oral Pathol* 29:603-612, 1970.
5. Walton RE, Torabinejad M: Obturación. In *Endodoncia*, Walton RE, Torabinejad M, Eds. 1st ed. Mexico: Nueva Editorial Interamericana SA, pp 241-268, 1990.
6. Coodlidge ED: Reaction of dog tissue to drugs used in root canal treatment. *J Am Dental Assoc* 19:747-59, 1932.
7. Seltzer S, Bender IB. Chemical irritants. In *The Dental Pulp. Biologic Considerations in Dental Procedures*, Seltzer S, Bender IB, Eds. 3rd ed. St. Louis: Ishiyaku Euroamerica, Inc., pp 215-37, 1990.
8. Myers DR, Pashley DH, Whitford GM, McKinney RV: Tissue changes induced by the absorption of formocresol from pulpotomy sites in dogs. *Pediatr Dent* 5:6-8, 1983.
9. Straffon LH, Han SS: Effect of varying concentrations of formocresol on RNA synthesis of connective tissues in sponge implants. *Oral Surg Oral Med Oral Pathol* 29:915-25, 1970.
10. Cunningham KW, Lazzeri EP, Ranly DM: The effect of formocresol and glutaraldehyde on certain enzymes in bovine dental pulp. *Oral Surg Oral Med Oral Pathol* 54:100-103, 1982.
11. Langeland K, Dowden WE, Tronstad L, Langeland LK: Human pulp changes of iatrogenic origin. *Oral Surg Oral Med Oral Pathol* 32:943-49, 1971.
12. Rölling I, Hasselgren G, Tronstad L: Morphologic and enzyme histochemical observations on the pulp of human primary molars 3 to 5 years after formocresol treatment. *Oral Surg Oral Med Oral Pathol* 42:518-28, 1976.
13. Pashley EL, Myers DR, Pashley DH, Whitford GM: Systemic distribution of 14C-formaldehyde from formocresol-treated pulpotomy sites. *J Dent Res* 59:602-608, 1980.
14. Osetek EM: Endodontic medicaments and irrigation solutions. In *Clinical Pharmacology in Dental Practice*, Holroyd SV, Wynn RL, Requa-Clark B, Eds. 4th ed. St. Louis: CV Mosby Co, pp 505-519, 1988.
15. Jendresen MD, Phillips RW: A comparative study of four zinc oxide and eugenol formulations as restorative materials. II. *J Prosthet Dent* 21:300-308, 1969.
16. Brännström M, Nordenvall KJ, Torstenson B: Pulpal reaction to IRM cement: an intermediate restorative material containing eugenol. *ASDC J Dent Child* 48:259-63, 1981.
17. Glass RL, Zander HA: Pulp healing. *J Dent Res* 28:97-107, 1949.
18. Brännström M, Nyborg H: Pulp reaction to a temporary zinc oxide/eugenol cement. *J Prosthet Dent* 35:185-91, 1976.
19. Webb JG Jr, Bussell NE: A comparison of the inflammatory response produced by commercial eugenol and purified eugenol. *J Dent Res* 60:1724-28, 1981.
20. Toriya Y, Hashiguchi I, Maeda K: Immunohistochemical examination of the distribution of macrophages and CGRP-immunoreactive nerve fibers in induced rat periapical lesions. *Endod Dent Traumatol* 13:6-12, 1997.
21. Kawashima N, Okiji T, Kosaka T, Suda H: Kinetics of macrophages and lymphoid cells during development of experimentally induced periapical lesions in rat molars: a quantitative immunohistochemical study. *J Endod* 22:311-16, 1996.

22. Marton IJ, Kiss C: Characterization of inflammatory cell infiltrate in dental periapical lesions. *Int Endod J* 26:131–36, 1993.
23. Trowbridge HO: Immunological aspects of chronic inflammation and repair. *J Endod* 16:54–61, 1990.
24. Unanue ER, Allen PM: The basis for the immunoregulatory role of macrophages and other accessory cells. *Science* 236:551–57, 1987.
25. Male D, Champion B, Cooke A: *Advanced Immunology*, 2nd ed. Philadelphia:JB Lippincott, 5.1–15.12, 1989.
26. Segura JJ, Guerrero JM, López-González MA, Calvo JR: Vasoactive intestinal peptide (VIP) inhibits substrate adherence capacity of rat peritoneal macrophages by a mechanism that involves cAMP. *Cell Adhes Commun* 1:213–21, 1993.
27. De la Fuente M, Del Río M, Ferrandez MD, Hernanz A: Modulation of phagocytic function in murine peritoneal macrophages by bombesin, gastrin-releasing peptide and neuromedin C. *Immunology* 73:205–211, 1991.
28. Noga SJ, Normann SJ, Weiner RS: Methods in laboratory investigation: isolation of guinea pig monocytes and Kurloff cells: characterization of monocyte subsets by morphology, cytochemistry, and adherence. *Lab Invest* 51:244–52, 1984.
29. Beekhuizen H, Corsèl-van Tilburg AJ, van Furth R: Characterization of monocyte adherence to human macrovascular and microvascular endothelial cells. *J Immunol* 145:510–18, 1990.
30. Garrido JJ, Arahuetes RM, Hernanz A, De la Fuente M: Modulation by neurotensin and neuromedin N of adherence and chemotaxis capacity of murine lymphocytes. *Regul Pept* 41:27–37, 1992.
31. Segura JJ, Llamas R, Jiménez Rubio-Manzanares A, Jiménez-Planas A, Guerrero JM, Calvo JR: Calcium hydroxide inhibits substrate adherence capacity of macrophages. *J Endodon*, in press.
32. Rotberg SJ, De Shazer DO: The complexing action of eugenol on sound dentin. *J Dent Res* 45:307–310, 1966.
33. Segura JJ, Calvo JR, Guerrero JM, Sampedro C, Jimenez A, Llamas R: Te disodium salt of EDTA inhibits the binding of vasoactive intestinal peptide to macrophage membranes: endodontic implications. *J Endod* 22:337–40.

## ***Pediatric Dentistry can communicate on line!***

Please direct questions, comments, or letters to the editor to:

**Dr. Milton I. Houpt, Editor-in-Chief**  
**houpt@umdnj.edu** (*Internet*)

You may also send correspondence or questions about manuscript preparation or status to:

**Diana Dodge, Journal Editor**  
**ddodge@aapd.org** (*Internet*)

We welcome your comments and questions. However, please follow the Instructions to Contributors on the web at <http://aapd.org> for manuscript submission procedures.

